


IN THE CLAIMS:

Please amend claims 1-54 as follows:



1. (original) A compound of formula I:



I

or a pharmaceutically acceptable salt, or mixtures thereof,  
wherein:

W is:



wherein each R<sub>6</sub> is independently:

- hydrogen-,
- (C1-C12)-aliphatic-,
- (C6-C10)-aryl-,
- (C6-C10)-aryl-(C1-C12)aliphatic-,
- (C3-C10)-cycloalkyl- or cycloalkenyl-,
- [(C3-C10)-cycloalkyl- or cycloalkenyl]-(C1-C12)-aliphatic-,
- (C3-C10)-heterocyclyl-,
- (C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,
- (C5-C10)-heteroaryl-, or
- (C5-C10)-heteroaryl-(C1-C12)-aliphatic-, or

wherein up to 3 aliphatic carbon atoms in each  $R_6$  may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)- in a chemically stable arrangement;

wherein  $R_6$  may be optionally substituted with up to 3 J substituents; or

two  $R_6$  groups, together with the nitrogen atom to which they are bound, may optionally form a 5- to 6-membered aromatic or a 3- to 7-membered saturated or partially unsaturated ring system wherein up to 3 ring atoms may be optionally replaced with N, NH, O, S, SO, and SO<sub>2</sub>, wherein said ring system may be optionally fused to a (C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or a (C3-C10)heterocyclyl, wherein any ring has up to 3 substituents selected independently from J;

wherein each  $R_8$  is independently -OR'; or the  $R_8$  groups together with the boron atom, may optionally form a (C3-C10)-membered heterocyclic ring wherein each  $R_8$  is independently -OR'; or the  $R_8$  groups together with the boron atom, may optionally form a (C3-C10)-membered heterocyclic ring having, in addition to the boron, up to 3 ring atoms optionally replaced with N, NH, O, S, SO, and SO<sub>2</sub>;

J is halogen, -OR', -NO<sub>2</sub>, -CN, -CF<sub>3</sub>, -OCF<sub>3</sub>, -R', oxo, thioxo, =N(R'), =N(OR'), 1,2-methylenedioxy, 1,2-ethylenedioxy, -N(R')<sub>2</sub>, -SR', -SOR', -SO<sub>2</sub>R', -SO<sub>2</sub>N(R')<sub>2</sub>, -SO<sub>3</sub>R', -C(O)R', -C(O)C(O)R', -C(O)C(O)OR', -C(O)C(O)NR', -C(O)CH<sub>2</sub>C(O)R', -C(S)R', -C(S)OR', -C(O)OR', -OC(O)R', -C(O)N(R')<sub>2</sub>, -OC(O)N(R')<sub>2</sub>, -C(S)N(R')<sub>2</sub>, -(CH<sub>2</sub>)<sub>0-2</sub>NHC(O)R', -N(R')N(R')COR', -N(R')N(R')C(O)OR', -N(R')N(R')CON(R')<sub>2</sub>, -N(R')SO<sub>2</sub>R', -N(R')SO<sub>2</sub>N(R')<sub>2</sub>, -N(R')C(O)OR', -N(R')C(O)R', -N(R')C(S)R', -N(R')C(O)N(R')<sub>2</sub>, -N(R')C(S)N(R')<sub>2</sub>, -N(COR')COR', -N(OR')R', -C(=NH)N(R')<sub>2</sub>, -C(O)N(OR')R',

-C(=NOR')R', -OP(O)(OR')<sub>2</sub>, -P(O)(R')<sub>2</sub>, -P(O)(OR')<sub>2</sub>, or  
-P(O)(H)(OR'); wherein;

R' is independently selected from:

hydrogen-,  
(C1-C12)-aliphatic-,  
(C3-C10)-cycloalkyl- or -cycloalkenyl-,  
[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C6-C10)-aryl-,  
(C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-heterocyclyl-,  
(C3-C10)-heterocyclyl-(C1-C12)aliphatic-,  
(C5-C10)-heteroaryl-, and  
(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein up to 5 atoms in R' may be optionally and  
independently substituted with J;

wherein two R' groups bound to the same atom may  
optionally form a 5- to 6-membered aromatic or a 3- to  
7-membered saturated or partially unsaturated ring  
system wherein up to 3 ring atoms may be optionally  
replaced with a heteroatom independently selected from  
N, NH, O, S, SO, and SO<sub>2</sub>, wherein said ring system may  
be optionally fused to a (C6-C10)aryl,  
(C5-C10)heteroaryl, (C3-C10)cycloalkyl, or a  
(C3-C10)heterocyclyl, wherein any ring has up to 3  
substituents selected independently from J;

R<sub>5</sub> and R<sub>5</sub>' are each independently hydrogen or (C1-C12)-  
aliphatic, wherein any hydrogen may be optionally  
replaced with halogen; wherein any terminal carbon atom  
of R<sub>5</sub> may be optionally substituted with sulfhydryl or  
hydroxy; or R<sub>5</sub> is Ph or -CH<sub>2</sub>Ph and R<sub>5</sub>' is H, wherein said  
Ph or -CH<sub>2</sub>Ph group may be optionally substituted with up  
to 3 substituents independently selected from J; or

R<sub>5</sub> and R<sub>5'</sub> together with the atom to which they are bound may optionally form a 3- to 6-membered saturated or partially unsaturated ring system wherein up to 2 ring atoms may be optionally replaced with N, NH, O, SO, or SO<sub>2</sub>; wherein said ring system has up to 2 substituents selected independently from J;

R<sub>2</sub>, R<sub>4</sub>, and R<sub>7</sub> are each independently:

hydrogen-,

(C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl-(C1-C12)-aliphatic-, or

(C6-C10)-aryl-(C1-C12)-aliphatic-;

wherein up to two aliphatic carbon atoms in each of R<sub>2</sub>, R<sub>4</sub>, and R<sub>7</sub> may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)- in a chemically stable arrangement;

wherein each of R<sub>2</sub>, R<sub>4</sub>, and R<sub>7</sub> may be independently and optionally substituted with up to 3 substituents independently selected from J;

R<sub>1</sub> and R<sub>3</sub> are each independently:

(C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

[(C3-C10)-cycloalkyl- or -cycloalkenyl]-(C1-C12)-aliphatic-,

(C6-C10)-aryl-(C1-C12)aliphatic-, or

(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein up to 3 aliphatic carbon atoms in each of R<sub>1</sub> and R<sub>3</sub> may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)- in a chemically stable arrangement;

wherein each of R<sub>1</sub> and R<sub>3</sub> may be independently and optionally substituted with up to 3 substituents independently selected from J;

R<sub>9</sub>, R<sub>9'</sub>, R<sub>10</sub>, and R<sub>10'</sub> are each independently -X-Y-Z;

X is a bond, -C(H)(R<sub>6</sub>)-, -O-, -S-, or -N(R<sub>11</sub>)-;

R<sub>11</sub> is:

hydrogen-,  
(C1-C12)-aliphatic-,  
(C6-C10)-aryl-,  
(C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-cycloalkyl- or cycloalkenyl-,  
[(C3-C10)-cycloalkyl- or cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C3-C10)-heterocyclyl-,  
(C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,  
(C5-C10)-heteroaryl-, or  
(C5-C10)-heteroaryl-(C1-C12)-aliphatic-,

wherein up to 3 aliphatic carbon atoms in each  $R_{11}$   
may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-,  
-O-, -N-, or -N(H)- in a chemically stable arrangement;

wherein  $R_{11}$  may be optionally substituted with up to  
3 J substituents; or

wherein  $R_{11}$  and Z together with the atoms to which  
they are bound, optionally form a nitrogen containing  
5-7-membered mono- or 6-11-membered bicyclic ring  
system optionally substituted with up to 3 J  
substitutents, wherein up to 3 ring atoms in said ring  
system may be optionally replaced with O, NH, S, SO, or  
SO<sub>2</sub> in a chemically stable arrangement;

Y is a bond, -CH<sub>2</sub>-, -C(O)-, -C(O)C(O)-, -S(O)-, S(O)<sub>2</sub>-, or -  
S(O)(NR<sub>12</sub>)-;

$R_{12}$  is:

hydrogen-,  
(C1-C12)-aliphatic-,  
(C6-C10)-aryl-,  
(C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-cycloalkyl- or cycloalkenyl-,  
[(C3-C10)-cycloalkyl- or cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C3-C10)-heterocyclyl-,

(C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,  
(C5-C10)-heteroaryl-, or  
(C5-C10)-heteroaryl-(C1-C12)-aliphatic-,

wherein up to 3 aliphatic carbon atoms in each R<sub>12</sub>  
may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-,  
-O-, -N-, or -N(H)-, in a chemically stable  
arrangement;

wherein R<sub>12</sub> may be optionally substituted with up to  
3 J substituents;

Z is:

hydrogen-,  
(C1-C12)-aliphatic-,  
(C3-C10)-cycloalkyl- or -cycloalkenyl-,  
[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C6-C10)-aryl-,  
(C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-heterocyclyl-,  
(C3-C10)-heterocyclyl-(C1-C12)aliphatic-,  
(C5-C10)-heteroaryl-, or  
(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein up to three aliphatic carbon atoms in Z may be  
optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or  
-N(H)-, in a chemically stable arrangement;

wherein any ring may be optionally fused to a  
(C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or  
(C3-C10)heterocyclyl;

wherein Z may be independently and optionally  
substituted with up to 3 substituents independently  
selected from J;

V is -C(O)-, -S(O)-, or -S(O)<sub>2</sub>-;

R is -C(O)-, -S(O)-, -S(O)<sub>2</sub>-, -N(R<sub>12</sub>)-, -O-, or a bond;

T is:

(C1-C12)-aliphatic-;

(C6-C10)-aryl-,  
(C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-cycloalkyl or -cycloalkenyl-,  
[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C3-C10)-heterocyclyl-,  
(C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,  
(C5-C10)-heteroaryl-, or  
(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein up to 3 aliphatic carbon atoms in T may be replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)-, in a chemically stable arrangement;

wherein each T may be optionally substituted with up to 3 J substituents; or

T is selected from -N(R<sub>6</sub>)(R<sub>6</sub>); and

R<sub>6</sub> is

hydrogen-,  
(C1-C12)-aliphatic-,  
(C6-C10)-aryl-,  
(C6-C10)-aryl-(C1-C12)aliphatic-,  
(C3-C10)-cycloalkyl- or cycloalkenyl-,  
[(C3-C10)-cycloalkyl- or cycloalkenyl]-(C1-C12)-  
aliphatic-,  
(C3-C10)-heterocyclyl-,  
(C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,  
(C5-C10)-heteroaryl-, or  
(C5-C10)-heteroaryl-(C1-C12)-aliphatic-, or



wherein up to 3 aliphatic carbon atoms in each R<sub>6</sub> may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)- in a chemically stable arrangement;

wherein R<sub>6</sub> may be optionally substituted with up to 3 J substituents; or

R<sub>6</sub> and R<sub>6</sub>, together with the nitrogen atom to which they are bound, may optionally form a (C3-C10)-

heterocyclic ring system wherein said ring system may be optionally substituted with up to 3 substituents independently selected from J.

2. (original) The compound according to claim 1, wherein the



in R<sub>9</sub>, R<sub>10</sub>, and R<sub>10'</sub>, X and Y are both a bond and Z is hydrogen; and in R<sub>9'</sub>;

X is a bond;

Y is a bond, -CH<sub>2</sub>-, or -C(O)-; and

Z is (C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-aliphatic-,

(C6-C10)-aryl-,

(C6-C10)-aryl-(C1-C12)aliphatic-,

(C3-C10)-heterocyclyl-,

(C3-C10)-heterocyclyl-(C1-C12)aliphatic-,

(C5-C10)-heteroaryl-, or

(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein up to three aliphatic carbon atoms in Z may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)-, in a chemically stable arrangement;

wherein any ring may be optionally fused to a (C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or (C3-C10)heterocyclyl;



wherein Z may be independently and optionally substituted with up to 3 substituents independently selected from J.

3. (original) The compound according to claim 2, wherein in R<sub>9</sub>;

X is a bond;

Y is a bond; and

Z is (C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-aliphatic-,

(C6-C10)-aryl-,

(C6-C10)-aryl-(C1-C12)aliphatic-,

(C5-C10)-heteroaryl-, or

(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein up to three aliphatic carbon atoms in Z may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)-, in a chemically stable arrangement;

wherein any ring may be optionally fused to a (C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or (C3-C10)heterocyclyl;

wherein Z may be independently and optionally substituted with up to 3 substituents independently selected from J.

4. (original) The compound according to claim 3, wherein in

R<sub>9</sub>;

X is a bond;

Y is a bond; and

Z is (C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,


[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
aliphatic-, or

(C6-C10)-aryl-(C1-C12)aliphatic-,


wherein up to three aliphatic carbon atoms in Z may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)-, in a chemically stable arrangement;

wherein Z may be independently and optionally substituted with up to 3 substituents independently selected from J.

5. (original) The compound according to claim 4, wherein R<sub>9</sub> is



6. (original) The compound according to claim 5, wherein R<sub>9</sub> is



7. (original) The compound according to claim 6, wherein R<sub>9</sub> is ethyl.

8. (original) The compound according to claim 1, wherein in  $R_9$ ,  $R_{10}$ , and  $R_{10'}$ , X and Y are both a bond and Z is hydrogen; and in  $R_9$ ;  
X is a bond;  
Y is  $-C(O)-$ ; and  
Z is (C1-C12)-aliphatic-, or  
(C3-C10)-heterocyclyl-(C1-C12)aliphatic-;  
wherein up to three aliphatic carbon atoms in Z may be optionally replaced with S,  $-S(O)-$ ,  $-S(O)_2-$ ,  $-O-$ ,  $-N-$ , or  $-N(H)-$ , in a chemically stable arrangement;  
wherein any ring may be optionally fused to a (C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or (C3-C10)heterocyclyl;  
wherein Z may be independently and optionally substituted with up to 3 substituents independently selected from J.

9. (original) The compound according to claim 8, wherein Z is  $-O-(C1-C6)-$ aliphatic or  $-N(R')_2$ , wherein the two  $R'$  groups bound to the nitrogen atom may optionally form a 3- to 7-membered saturated or partially unsaturated ring system wherein up to 3 ring atoms may be optionally replaced with a heteroatom independently selected from N, NH, O, S, SO, and  $SO_2$ , wherein said ring system may be optionally fused to a (C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or a (C3-C10)heterocyclyl, wherein any ring has up to 3 substituents selected independently from J.

10. (original) The compound according to claim 8, wherein Z is  $-N(R')_2$ , wherein the two  $R'$  groups bound to the nitrogen atom may optionally form a 3- to 7-membered saturated or partially unsaturated ring system wherein up to 3 ring atoms may be optionally replaced with a heteroatom

independently selected from N, NH, O, S, SO, and SO<sub>2</sub>, wherein said ring system may be optionally fused to a (C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or a (C3-C10)heterocyclyl, wherein any ring has up to 3 substituents selected independently from J.

11. (original) The compound according to claim 1, wherein in

R<sub>9</sub>, and R<sub>10</sub>, X and Y are a bond and Z is hydrogen; and in each of R<sub>9</sub>, and R<sub>10</sub>, independently;

X is a bond;

Y is a bond; and

Z is (C1-C12)-aliphatic-,

(C3-C10)-cycloalkyl- or -cycloalkenyl-,

[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-aliphatic-,

(C6-C10)-aryl-,

(C6-C10)-aryl-(C1-C12)aliphatic-,

(C3-C10)-heterocyclyl-,

(C3-C10)-heterocyclyl-(C1-C12)aliphatic-,

(C5-C10)-heteroaryl-, or

(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein up to three aliphatic carbon atoms in Z may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)-, in a chemically stable arrangement;

wherein any ring may be optionally fused to a (C6-C10)aryl, (C5-C10)heteroaryl, (C3-C10)cycloalkyl, or (C3-C10)heterocyclyl;

wherein Z may be independently and optionally substituted with up to 3 substituents independently selected from J.

12. (original) The compound according to claim 11, wherein Z, in each of R<sub>9</sub>, and R<sub>10</sub>, independently, is

(C1-C12)-aliphatic-,  
(C3-C10)-cycloalkyl- or -cycloalkenyl-, or  
[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-  
aliphatic-;

wherein up to three aliphatic carbon atoms in Z may be optionally replaced with S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)-, in a chemically stable arrangement;


wherein Z may be independently and optionally substituted with up to 3 substituents independently selected from J.

13. (original) The compound according to claim 12, wherein Z, in each of R<sub>9</sub> and R<sub>10</sub>, independently, is (C1-C6)-aliphatic-.

14. (original) The compound according to claim 1, wherein in R<sub>10</sub> and R<sub>10'</sub>, X and Y are a bond and Z is hydrogen; and in each of R<sub>9</sub> and R<sub>9'</sub>,  
X is a bond,  
Y is a bond, and  
Z is (C1-C6)-aliphatic-,


wherein Z may be independently and optionally substituted with up to 3 substituents independently selected from J.

15. (original) The compound according to any one of claims 1-14, wherein W is:




wherein in the W, the  $\text{NR}_6\text{R}_6$  is selected from  $-\text{NH}-(\text{C1-C6 aliphatic})$ ,  $-\text{NH}-(\text{C3-C6 cycloalkyl})$ ,  $-\text{NH}-\text{CH}(\text{CH}_3)-\text{aryl}$ , or  $-\text{NH}-\text{CH}(\text{CH}_3)-\text{heteroaryl}$ , wherein said aryl or said heteroaryl is optionally substituted with up to 3 halogens.


16. (original) The compound according to claim 15, wherein in the W, the  $\text{NR}_6\text{R}_6$  is:




17. (original) The compound according to claim 16, wherein in the W, the  $\text{NR}_6\text{R}_6$  is:




18. (original) The compound according to claim 17, wherein in the W, the  $\text{NR}_6\text{R}_6$  is:




19. (original) The compound according to claim 18, wherein in the W, the  $\text{NR}_6\text{R}_6$  is:



20. (currently amended) The compound according to ~~any~~  
~~one of claims 1-19~~, wherein R<sub>5</sub> is hydrogen and R<sub>5</sub> is:




21. (original) The compound according to claim 20, wherein R<sub>5</sub> is hydrogen and R<sub>6</sub> is:




22. (currently amended) The compound according to ~~any~~  
~~one of~~ claims 1-21, wherein R<sub>2</sub>, R<sub>4</sub>, and R<sub>7</sub> are each  
independently H, methyl, ethyl, or propyl.

23. (original) The compound according to claim 22, wherein R<sub>2</sub>, R<sub>4</sub>, and R<sub>7</sub> are each hydrogen.


24. (currently amended) The compound according to ~~any~~  
~~one of claims 1-33~~, wherein R<sub>3</sub> is:




25. (original) The compound according to claim 24,  
wherein  $R_3$  is:




26. (original) The compound according to claim 25,  
wherein  $R_3$  is:



27. (currently amended) The compound according to ~~any~~  
~~one of~~ claims 1—26, wherein  $R_1$  is:




28. (original) The compound according to claim 27,  
wherein  $R_1$  is:




29. (original) The compound according to claim 18,  
wherein  $R_1$  is isopropyl or cyclohexyl.



30. (original) The compound according to claim 1, wherein the




radical is:



wherein:

$R_6$ ,  $R_{6'}$ ,  $R_7$ , and  $R_{12}$ , are as defined in claim 1.

31. (original) The compound according to claim 30, wherein in the



radical;

$R_{6'}$  and  $R_7$  are both hydrogen;

$R_6$  is:

(C1-C12)-aliphatic-;

(C6-C10)-aryl-,

(C6-C10)-aryl-(C1-C12)aliphatic-,

(C3-C10)-cycloalkyl or -cycloalkenyl-,

[(C3-C10)-cycloalkyl or -cycloalkenyl]-(C1-C12)-aliphatic-,

(C3-C10)-heterocyclyl-,

(C3-C10)-heterocyclyl-(C1-C12)-aliphatic-,

(C5-C10)-heteroaryl-, or

(C5-C10)-heteroaryl-(C1-C12)-aliphatic-;

wherein up to 3 aliphatic carbon atoms in  $R_6$  may be optionally replaced by S, -S(O)-, -S(O)<sub>2</sub>-, -O-,

-N-, or -N(H)-, in a chemically stable arrangement;  
and

wherein  $R_6$  may be optionally substituted with up to 3 substituents independently selected from J; and  $R_{12}$  is as defined in claim 1.

32. (original) The compound according to claim 31, wherein;

$R_6$  is:

(C1-C12)-aliphatic-;


(C6-C10)-aryl-(C1-C12)aliphatic-, or

(C3-C10)-cycloalkyl or -cycloalkenyl-;


wherein up to 3 aliphatic carbon atoms in  $R_6$  may be optionally replaced by S, -S(O)-, -S(O)<sub>2</sub>-, -O-, -N-, or -N(H)-, in a chemically stable arrangement;

wherein  $R_6$  may be optionally substituted with up to 3 substituents independently selected from J; and  $R_{12}$  is as defined in claim 1.



33. (original) The compound according to claim 32, wherein the radical is:




34. (original) The compound according to claim 33, wherein the radical is:



radical is:



or



35. (currently amended) The compound according to ~~any~~  
~~one of~~ claims 1-29, wherein;  
V is -C(O)-; and  
R is a bond.


36. (currently amended) The compound according to ~~any~~  
~~one of~~ claims 1-29, wherein;  
V is -C(O)-;  
R is a bond; and  
T is:

(C3-C10)-heterocyclyl- or (C5-C10)heteroaryl-;

wherein each T is optionally substituted with up to 3 J  
substituents.

37. (original) The compound according to claim 36,  
wherein T is (C5-C6)heterocyclyl- or (C5-C6)heteroaryl-;  
wherein each T is optionally substituted with up to 3 J  
substituents.


38. (original) The compound according to claim 37,  
wherein T is:











wherein:










Z' is independently O, S, NR', or C(R')<sub>2</sub>.











39. (original) The compound according to claim 38, wherein T is:













40. (original) The compound according to claim 1, wherein the compound is:










1	
2	
3	
4	
5	
6	
7	
8	











9	
10	
11	
12	
13	
14	
15	
16	
17	










18	
19	
20	
21	
22	
23	
24	
25	
26	
27	










28	
29	
30	
31	
32	
33	
34	
35	
36	
37	




38	
39	
40	
41	
42	
43	
44	
45	
46	

47	
48	
49	
50	
51	
52	
53	
54	
55	
56	

57	
58	
59	
60	
61	
62	
63	
64	
65	

66	
67	
68	
69	
70	
71	
72	
73	
74	



41. (currently amended) A pharmaceutical composition comprising a compound according to ~~any one of claims 1-40~~ or a pharmaceutically acceptable salt or mixtures thereof in an amount effective to inhibit a serine protease; and a acceptable carrier, adjuvant or vehicle.

42. (original) The composition according to claim 41, wherein said composition is formulated for administration to a patient.

43. (original) The composition according to claim 42, wherein said composition comprises an additional agent selected from an immunomodulatory agent; an antiviral agent; a second inhibitor of HCV protease; an inhibitor of another target in the HCV life cycle; and a cytochrome P-450 inhibitor; or combinations thereof.

44. (original) The composition according to claim 41, wherein said immunomodulatory agent is  $\alpha$ -,  $\beta$ -, or  $\gamma$ -interferon or thymosin; said antiviral agent is ribavirin, amantadine, or telbivudine; or said inhibitor of another

target in the HCV life cycle is an inhibitor of HCV helicase, polymerase, or metalloprotease.

45. (original) The composition according to claim 43, wherein said cytochrome P-450 inhibitor is ritonavir.

46. (currently amended) A method of inhibiting the activity of a serine protease comprising the step of contacting said serine protease with a compound according to ~~any one of~~ claims 1-40.

47. (original) The method according to claim 46, wherein said serine protease is an HCV NS3 protease.

48. (original) A method of treating an HCV infection in a patient comprising the step of administering to said patient a composition according to claim 42.

49. (original) The method according to claim 48, comprising the additional step of administering to said patient an additional agent selected from an immunomodulatory agent; an antiviral agent; a second inhibitor of HCV protease; an inhibitor of another target in the HCV life cycle; or combinations thereof; wherein said additional agent is administered to said patient as part of said composition according to claim 42 or as a separate dosage form.

50. (original) The method according to claim 49, wherein said immunomodulatory agent is  $\alpha$ -,  $\beta$ -, or  $\gamma$ -interferon or thymosin; said antiviral agent is ribavirin or amantadine; or said inhibitor of another target in the HCV life cycle is an inhibitor of HCV helicase, polymerase, or metalloprotease.

512. (currently amended) A method of eliminating or reducing HCV contamination of a biological sample or medical or laboratory equipment, comprising the step of contacting said biological sample or medical or laboratory equipment with a composition according to claim 41.

523. (currently amended) The method according to claim 512, wherein said sample or equipment is selected from blood, other body fluids, biological tissue, a surgical instrument, a surgical garment, a laboratory instrument, a laboratory garment, a blood or other body fluid collection apparatus; a blood or other body fluid storage material.

534. (currently amended) The method according to claim 523, wherein said body fluid is blood.